

REGULARIZATION AND STABILITY OF SOLUTIONS OF SYSTEM OF LINEAR INTEGRAL FREDHOLM EQUATIONS OF THE FIRST KIND ON A SEMI-AXIS

Asanov A.¹ , Kadenova Z.A.², Bekeshova D.A.³,

¹*Department of Mathematics , Kyrgyz-Turkish Manas University*

^{2,3}*Institute of Mathematics of NAS of the Kyrgyz Republic*

In this article, regularizations and stability of solutions of system linear integral Fredholm equations of the first kind are obtained by the methods of functional analysis.

Key words: linear, inteqral equations, first kind, regularization.

Бул макалада функционалдык анализдин усулдарынын жардамы менен Фредгольмдун биринчи түрдөгү сызыктуу интегралдык тендемелер системасынын чечимдеринин регуляризациясы жана туруктуулугу алынды.

Урунттуу сөздөр: сызыктуу, интегралдык тендемелер, биринчи түрдөгү, регуляризация.

В данной работе, с помощью методов функционального анализа получены регуляризации и устойчивости решений систем линейных интегральных уравнений Фредгольма первого рода.

Ключевые слова: линейные, интегральные уравнения, первого рода, регуляризация.

1. Introduction

Inverse and ill-posed problems are currently attracting great interest. The theory and numerical methods for solving inverse and ill-posed problems were studied in [1-18]. The notion of correctness in the works of A.N. Tikhonov [1], M.M. Lavrent'ev [2] and V.K. Ivanov [3], different from the classical one, provided a means for studying ill-posed problems and stimulated interest in integral equations with large applied value.

The fundamental results for Fredholm integral equations of the first kind were obtained by M.M. Lavrentiev in [4], [5], where the regularizing operators in the sense of M.M. Lavrentiev.

In [10]-[12], uniqueness theorems were proved and regularization operators in the sense of Lavrent'ev were constructed for systems of linear Volterra and Fredholm integral equations of the third kind.

In [14]-[16] problems of uniqueness and stability of solutions for linear Fredholm integral equations of the first kind were investigated.

In this work, we apply the method of integral transformation to prove regularizations and stability of solutions of system of linear integral Fredholm equations of the first kind in the semi-axis.

Consider of the system of Fredholm linear integral equations

$$Ku \equiv \int_{-\infty}^a K(t,s)u(s)ds = f(t), \quad t \in (-\infty, a] \quad (1)$$

where

$$\int_{-\infty}^a \int_{-\infty}^a |K(t,s)|^2 ds dt < \infty,$$

$$K(t,s) = \begin{cases} A(t,s), & -\infty < s \leq t \leq a, \\ B(t,s), & -\infty < t \leq s \leq a. \end{cases} \quad (2)$$

$$A(t,s) = a_{ij}(t,s) = \begin{pmatrix} a_{11}(t,s) & a_{12}(t,s) & \dots & a_{1n}(t,s) \\ a_{21}(t,s) & a_{22}(t,s) & \dots & a_{2n}(t,s) \\ \dots & \dots & \dots & \dots \\ a_{n1}(t,s) & a_{n2}(t,s) & \dots & a_{nn}(t,s) \end{pmatrix}, \quad B(t,s) = b_{ij}(t,s) = \begin{pmatrix} b_{11}(t,s) & b_{12}(t,s) & \dots & b_{1n}(t,s) \\ b_{21}(t,s) & b_{22}(t,s) & \dots & b_{2n}(t,s) \\ \dots & \dots & \dots & \dots \\ b_{n1}(t,s) & b_{n2}(t,s) & \dots & b_{nm}(t,s) \end{pmatrix}$$

$$f(t) = (f_i(t)) = (f_1(t), \dots, f_n(t))^T, \quad u(t) = (u_i(t)) = (u_1(t), \dots, u_n(t))^T.$$

$B(t,s)$ and $A(t,s)$ -given matrix functions, $f(t)$ - known vector function, $u(t)$

- unknown vector function.

$$\{(t,s) : -\infty < s \leq t \leq a\}, \{(t,s) : -\infty < t \leq s \leq a\}.$$

We will assume that

$$\|K(t,s)\| \in L_2(R \times R), \quad \|f(t)\| \in L_2(R)$$

System of equations (1) by virtue of relation (2) can be expressed as

$$\int_{-\infty}^t A(t,s)u(s)ds + \int_t^a B(t,s)u(s)ds = f(t) \quad (3)$$

Both parts of (3) are scalarly multiplied by the $u(t)$ -vector function and), integrating the results on $a \leq t < \infty$, we obtain

$$\begin{aligned} \int_{-\infty}^a \int_{-\infty}^t \langle A(t,s)u(s), u(t) \rangle ds dt + \int_{-\infty}^a \int_t^a \langle B(t,s)u(s), u(t) \rangle ds dt &= \int_{-\infty}^a \langle f(t), u(t) \rangle dt \\ \int_{-\infty}^a \int_{-\infty}^t \langle A(t,s)u(s), u(t) \rangle ds dt + \int_{-\infty}^a \int_s^a \langle B^*(s,t)u(s), u(t) \rangle dt ds &= \int_{-\infty}^a \langle f(t), u(t) \rangle dt, \end{aligned} \quad (4)$$

where $B^*(s, t)$ is the transposed matrix to the matrix $B(s, t)$.

Integrating by parts and using the Dirichlet formula we have

$$\begin{aligned} \int_{-\infty}^a \int_s^a B(s,t)u(t)u(s)dt ds &= \int_{-\infty}^a \left[\int_{-\infty}^t B(s,t)u(s)dt \right] u(t)dt. \\ \int_{-\infty}^a \int_s^a \langle B^*(s,t)u(s), u(t) \rangle dt ds &= \int_{-\infty}^a \int_{-\infty}^t \langle B^*(s,t)u(s), u(t) \rangle dt ds \\ \int_{-\infty}^a \int_{-\infty}^t \langle [A(t,s) + B^*(s,t)]u(s), u(t) \rangle ds dt &= \int_{-\infty}^a \langle f(t), u(t) \rangle dt \end{aligned} \quad (5)$$

Denote

$$H(t,s) = \frac{1}{2} (A(t,s) + B^*(s,t)) \quad (t,s) \in G = \{-\infty < s < t \leq a\}$$

Then from (5) we obtain

$$2 \int_{-\infty}^a \int_{-\infty}^t \langle H(t,s)u(s), u(t) \rangle ds dt = \int_{-\infty}^a \langle f(t), u(t) \rangle dt. \quad (6)$$

$$\int_{-\infty}^a \int_{-\infty}^t H^2(t,s) dt ds < +\infty.$$

We introduce a new matrix function $M(t,s)$ as follows

$$M(t,s) = \begin{cases} H(t,s), & -\infty < s \leq t \leq a, \\ H(s,t), & -\infty < t \leq s \leq a. \end{cases} \quad (7)$$

It's clear that $M(t,s) = M(s,t)$, $(t,s) \in (-\infty, a] \times (-\infty, a]$.

It is easy to verify the validity of the equality

$$\int_{-\infty}^a \int_{-\infty}^a |M(t,s)|^2 ds dt < +\infty.$$

Then, it is known that

$$M(t,s) = \sum_{i=1}^{\infty} \frac{\varphi_i(t)\varphi_i(s)}{\lambda_i}, \quad (8)$$

where, λ_i - the characteristic numbers of the matrix kernel $M(t,s)$, which are arranged in ascending order of their modules, $|\lambda_1| \leq |\lambda_2| \leq \dots$ and $\varphi_1(t), \varphi_2(t) \dots$

the corresponding orthonormal eigen vector-functions.

It is assumed that $M(t,s)$ is a complete kernel and $0 < \lambda_1 \leq \lambda_2 \leq \dots$. In this case, the solution of equation (1) will be unique in $L_2(-\infty, a]$.

In what follows, we will assume that all characteristic numbers of the matrix kernel are positive.

For $u(t) = (u_i(t) \in L_2(R; R^n))$ we define the norm

$$\|u(t)\|_2 = \left(\sum_{i=1}^n \int_{-\infty}^a |u_i(t)|^2 dt \right)^{\frac{1}{2}} = \left(\int_{-\infty}^a \|u(t)\|^2 dt \right)^{\frac{1}{2}} = \left(\sum_{v=1}^{\infty} |u^{(v)}|^2 \right)^{\frac{1}{2}},$$

were

$$u^{(v)} = \int_{-\infty}^a \langle u(t), \varphi^{(v)}(t) \rangle dt = \int_{-\infty}^a \left(\sum_{i=1}^n u_i(t) \varphi_i^{(v)}(t) \right) dt, \quad (v=1,2,\dots).$$

We distinguish the family of correctness sets depending on the parameter as follows:

$$N_\alpha = \left\{ u(t) \in L_2((-\infty, a]; R^n) : \sum_{v=1}^{\infty} \lambda_v^{-\alpha} |u^{(v)}|^2 \leq c \right\}, \quad \text{where } c > 0, 0 < \alpha < \infty,$$

$$u^{(v)} = \int_{-\infty}^a \langle u(t), \varphi^{(v)}(t) \rangle dt, \quad (v = 1, 2, \dots). \quad (9)$$

Let $u(t) = (u_i(t)) \in N_\alpha$. Then

$$2 \int_{-\infty}^a \int_a^t \sum_{v=1}^{\infty} \lambda_v \left\langle \begin{pmatrix} \varphi_1^{(v)}(t) \\ \dots \\ \varphi_1^{(v)}(t) \end{pmatrix} (\varphi_1^{(v)}(s) \dots \varphi_n^{(v)}(s)) \begin{pmatrix} u_1(s) \\ \dots \\ u_n(s) \end{pmatrix}, u(t) \right\rangle ds dt = \int_{-\infty}^a \left[\sum_{i=1}^n f_i(t) u_i(t) \right] dt$$

$$\|u(t)\|_{L_2}^2 = \sum_{i=1}^{\infty} |u^{(v)}|^2 = \sum_{v=1}^{\infty} \lambda_v^\alpha \lambda_v^{-\alpha} |u^{(v)}|^2 = \lambda_1^\alpha \left(\sum_{v=1}^{\infty} \lambda_v^{-\alpha} |u^{(v)}|^2 \right) \leq c \lambda_1^\alpha,$$

$$\|u(t)\|_{L_2}^2 \leq c \lambda_1^\alpha. \quad (10)$$

We will assume that $f(t) \in K(N_\alpha)$. Then the system (1) has a solution $(u_i(t)) \in N_\alpha$ and by virtue of (6), (7) and (8) we have:

$$2 \int_{-\infty}^a \int_a^t \sum_{v=1}^{\infty} \lambda_v \left\langle \begin{pmatrix} \varphi_1^{(v)}(t) \\ \dots \\ \varphi_1^{(v)}(t) \end{pmatrix} (\varphi_1^{(v)}(s) \dots \varphi_n^{(v)}(s)) \begin{pmatrix} u_1(s) \\ \dots \\ u_n(s) \end{pmatrix}, u(t) \right\rangle ds dt = \int_{-\infty}^a \left[\sum_{i=1}^n f_i(t) u_i(t) \right] dt,$$

$$\sum_{v=1}^{\infty} 2 \int_{-\infty}^a \left[\int_a^t \langle u(s), \varphi^{(v)}(s) \rangle ds \right] \langle u(t), \varphi^{(v)}(t) \rangle dt = \int_{-\infty}^a \langle f(t), u(t) \rangle dt.$$

$$\sum_{v=1}^{\infty} \lambda_v \left| \int_{-\infty}^a \langle u(t), \varphi^{(v)}(t) \rangle dt \right|^2 = \int_{-\infty}^a \langle f(t), u(t) \rangle dt$$

$$\sum_{v=1}^{\infty} \lambda_v |u^{(v)}|^2 = \int_{-\infty}^a \langle f(t), u(t) \rangle dt$$

Hence, using Hölder's inequalities, we have

$$\sum_{v=1}^{\infty} \lambda_v |u^{(v)}| \leq \|f(t)\|_{L_2} \cdot \|u(t)\|_{L_2} \quad (11)$$

On the other side:

$$\|u(t)\|_{L_2}^2 = \sum_{v=1}^{\infty} \frac{|u^{(v)}|^{2\alpha}}{\lambda_v^{\frac{\alpha}{1+\alpha}}} \lambda_v^{\frac{\alpha}{1+\alpha}} |u^{(v)}|^{\frac{2}{1+\alpha}} \leq \left(\sum_{v=1}^{\infty} \frac{|u_v|^2}{\lambda_v} \right)^{\frac{\alpha}{1+\alpha}} \left(\sum_{v=1}^{\infty} \lambda_v^{-\alpha} |u^{(v)}|^2 \right)^{\frac{1}{1+\alpha}}$$

Here we have applied Hölder's inequality for $p = 1 + a, q = \frac{(1 + \alpha)}{\alpha}$. Taking into account $u(t) \in N_\alpha$ and (11), from the last inequality we have

$$\|u(t)\|_{L_2}^2 \leq c^{\frac{1}{1+\alpha}} (\|f(t)\|_{L_2} \|u(t)\|_{L_2})^{\frac{\alpha}{1+\alpha}}$$

Hence we obtain the following stability estimate

$$\|u(t)\|_{L_2} \leq c^{\frac{1}{2+\alpha}} \|f(t)\|_{L_2}^{\frac{\alpha}{2+\alpha}}, \quad 0 < \alpha < \infty. \quad (12)$$

Thus, the following theorem has been proved.

Theorem 1. Let the operator M generated by the matrix kernel $M(t, s)$ be positive, where it is defined $M(t, s)$ by formula (7). Then on the set $K(N_\alpha)$ ($K(N_\alpha)$ image when N_α displayed by the operator K) the operator K^{-1} , the

inverse of K , is uniformly continuous with the Hölder exponent $\frac{\alpha}{2+\alpha}$, i.e. estimate (12) is valid.

Let us show that the solution of the system of equations

$$\varepsilon u(t, \varepsilon) + \int_{-\infty}^a K(t, s) u(s, \varepsilon) ds = f(t), \quad t \in (-\infty, a], \varepsilon > 0 \quad (13)$$

will be regularizing for equation (1) on the set N_α .

Indeed, by making the following substitution into the in system equations (13)

$$u(t, \varepsilon) = u(t) + \xi(t, \varepsilon),$$

where $u(t) \in N_\alpha$ - solution of systems of equations (1), we obtain

$$\varepsilon \xi(t, \varepsilon) + \int_{-\infty}^a K(t, s) \xi(s, \varepsilon) ds = -\varepsilon u(t)$$

Scalarly multiplying the last system of equations by $\xi(t, \varepsilon)$ and integrating from $-\infty$ to a , taking into account (2) and (8), we have:

$$\varepsilon \|\xi(t, \varepsilon)\|^2 + \sum_{v=1}^{\infty} \lambda_v^{-1} |\xi_v(\varepsilon)|^2 \leq \varepsilon \sum_{v=1}^{\infty} |u^{(v)}| |\xi_v(\varepsilon)| \quad (14)$$

where $\xi_i(\varepsilon)$ -are the Fourier coefficients for the function $\xi(t, \varepsilon)$, according to the orthonormal system $\phi^{(v)}(t) = \{\phi_i^{(v)}(t)\}$ that is

$$\xi_v(\varepsilon) = \int_a^\infty \langle \xi(t, \varepsilon), \phi_v(t) \rangle dt, \quad (v = 1, 2, \dots)$$

Applying the Hölder inequality for $p = q = \frac{1}{2}$, from (14) we obtain

$$\|\xi(t, \varepsilon)\|_{L_2} \leq \|u(t)\|_{L_2} \quad (15)$$

$$\sum_{v=1}^{\infty} \lambda_v |\xi_v(\varepsilon)|^2 \leq \varepsilon \|u(t)\|_{L_2}^2 \leq \varepsilon c \lambda_1^\alpha, \quad \varepsilon > 0, \quad (16)$$

on the other side

$$\sum_{v=1}^{\infty} |u^{(v)}| |\xi_v(\varepsilon)| = \sum_{v=1}^{\infty} \frac{|\xi_v(\varepsilon)|^{\frac{\alpha}{1+\alpha}}}{\lambda_v^{\frac{\alpha}{2(1+\alpha)}}} \cdot \lambda_v^{\frac{\alpha}{2(1+\alpha)}} |u^{(v)}|^{\frac{1}{1+\alpha}} |\xi_v(t, \varepsilon)|^{\frac{1}{1+\alpha}} |u^{(v)}|^{\frac{\alpha}{1+\alpha}}$$

Therefore, after applying the generalized Hölder inequality to the right-hand side for

$$p = \frac{2(1+\alpha)}{\alpha}, \quad q = 2(1+\alpha), \quad m = 2(1+\alpha), \quad n = \frac{2(1+\alpha)}{\alpha} \quad \text{we have}$$

$$\sum_{v=1}^{\infty} |u^{(v)}| |\xi_v(\varepsilon)| \leq \left(\sum_{v=1}^{\infty} \lambda_v |\xi_v(\varepsilon)|^2 \right)^{\frac{1}{p}} \left(\sum_{v=1}^{\infty} \frac{|u^{(v)}|^2}{\lambda_v^\alpha} \right)^{\frac{1}{q}} \|\xi(t, \varepsilon)\|_{L_2}^{\frac{2}{q}} \|u(t)\|_{L_2}^{\frac{2}{p}}$$

$$\sum_{v=1}^{\infty} |u^{(v)}| |\xi_v(\varepsilon)| \leq \left(\sum_{v=1}^{\infty} |\xi_v(\varepsilon)|^2 \lambda_v^{-1} \right)^{\frac{1}{(1+\alpha)2}} \left(\sum_{v=1}^{\infty} \lambda_v^\alpha |u^{(v)}|^2 \right)^{\frac{1}{(1+\alpha)2}} \|\xi(t, \varepsilon)\|_{L_2}^{\frac{1}{1+\alpha}} \|u(t)\|_{L_2}^{\frac{\alpha}{1+\alpha}}$$

$$\sum_{v=1}^{\infty} |u^{(v)}| |\xi_v(\varepsilon)| \leq \left(\sum_{v=1}^{\infty} |\xi_v(\varepsilon)|^2 \lambda_v^{-1} \right)^{\frac{1}{p}} \left(\sum_{v=1}^{\infty} \lambda_v^\alpha |u^{(v)}|^2 \right)^{\frac{1}{q}} \|\xi(t, \varepsilon)\|_{L_2}^{\frac{2}{q}} \|u(t)\|_{L_2}^{\frac{2}{p}}$$

Further in force

$u(t) \in N_\alpha$, (15) and (16) from the last inequality we have

$$\sum_{v=1}^{\infty} \|u^{(v)}\|_{\xi_v}(\varepsilon) \leq (\varepsilon c \lambda_1^\alpha)^{\frac{1}{p}} c^{\frac{1}{q}} (c \lambda_1^\alpha)^{\frac{p+q}{pq}}.$$

Hence, substituting $p = \frac{2(1+\alpha)}{\alpha}$, $q = 2(1+\alpha)$, we get

$$\sum_{v=1}^{\infty} \|u^{(v)}\|_{\xi_v}(\varepsilon) \leq c^{\frac{1}{2(1+\alpha)}} (c \lambda_1^\alpha)^{\frac{1}{2}} (\varepsilon c \lambda_1^\alpha)^{\frac{\alpha}{2(1+\alpha)}}, \quad (17)$$

$$\sum_{v=1}^{\infty} \|u^{(v)}\|_{\xi_v}(\varepsilon) \leq c \lambda_1^{\frac{1}{2(1+\alpha)}} c^{\frac{1}{2}} c^{\frac{\alpha}{2(1+\alpha)}} \lambda_1^{\frac{\alpha}{2}} \lambda_1^{\frac{\alpha^2}{2(1+\alpha)}} \varepsilon^{\frac{\alpha}{2(1+\alpha)}}$$

$$\sum_{v=1}^{\infty} \|u^{(v)}\|_{\xi_v}(\varepsilon) \leq c \lambda_1^{\frac{\alpha(2\alpha+1)}{2(1+\alpha)}} \varepsilon^{\frac{\alpha}{2(1+\alpha)}} \quad (18)$$

Taking into account (18), from (14) we have

$$\|u(t, \varepsilon) - u(t)\|_{L_2} \leq c^{\frac{1}{2}} \lambda_1^{\frac{\alpha(2\alpha+1)}{4(1+\alpha)}} \varepsilon^{\frac{\alpha}{4(1+\alpha)}}, 0 < a < \infty \quad (19)$$

Thus proven.

Theorem 2. Let the operator M generated by the matrix kernel $M(t, s)$ be positive and $f(t) \in K(N_\alpha)$. Then estimate (19) is valid, where $u(t, \varepsilon)$ -is the solution of the system (13), $u(t)$ is the solution of the system (1) $M(t, s)$ and is determined by formula (8).

REFERENCES

1. Tikhonov A.N. On methods for solving ill-posed problems. // In the book: Abstracts of reports, International Congress of Mathematicians. - M., 1966. (russ).
2. Lavrent'ev M. M. On some ill-posed problems of mathematical physics. - Novosibirsk: Publishing House of Sib. Department of the Academy of Sciences of the USSR, 1962. - 92 p.(russ).
3. Ivanov V.K. About ill-posed tasks. Differential Equations.-1968.- No. 2.-P.61.
4. Lavrent'ev M. M. On integral equations of the first kind Doklady of the Academy of Sciences of the USSR. 1959. V. 127. No. 1. S. 31.(russ).
5. Lavrent'ev M. M., Romanov V. G., Shishatskii S. P. Ill-Posed Problems of Mathematical Physics and Analysis (Providence, RI: American Mathematical Society). – 1986.
6. Apartsyn, A. S. Nonclassical Linear Volterra Equations of the First Kind / A. S. Apartsyn. – Utrecht-Boston: VSP, 2003. – 168 p. – ISBN 90-6764-375-0. – EDN RUCKLT.
7. Magnitskii N. A. Linear Volterra Integral Equations of the First and Third Kind // Doklady Akademii Nauk. - 1991. - Vol. 317, No 2.-Pp. 330-333.
8. Bukhgeim A.L. Volterra Equations and Inverse Problems, VSP, Utrecht, The Netherlands, 1999.
9. A.Asanov, Regularization, Uniqueness and Existence of Solutions of Volterra Equations of the First Kind. VSP, Utrecht, p.272. (1998).
10. Imanaliev, M. I., Asanov A., Solutions of system of nonlinear Volterra integral equations of the first kind, Soviet Math. Dokl. 309, (1989) 1052–1055.
11. Imanaliev, M. I., Asanov A., Asanov R. A. On a class of systems of linear and nonlinear Fredholm integral equations of the third kind with multipoint singularities // Differential Equations. - 2018. - T. 54. - No. 3. - P. 387. – DOI 10.1134/S037406411803010X. – EDN YQYHAP. (Russ).
12. Imanaliev, M. I., Asanov A. Regularization and uniqueness of solutions to systems of nonlinear Volterra integral equations of the third kind // Doklady

- Mathematics. – 2007. – Vol. 415. – No 1. – P. 14-17. – DOI 10.1134/S1064562407040035. – EDN XLWHPB.(Russ)
13. Imanaliev M. I., Asanov A., Asanov R. A. A class of systems of linear Fredholm integral equations of the third kind // *Doklady Mathematics*. – 2011. – Vol. 83. – No 2. – P. 227-231. – DOI 10.1134/S1064562411020293. – EDN XKKSFX.
 14. Asanov A., Kadenova Z. A. Regularization and Stability of Systems of Linear Integral Fredholm Equations of the First Kind, *Vestn. Samar. Gos. Tekhn. Univ., Ser. Fiz.-Mat. Nauki [J. Samara State Tech. Univ., Ser. Phys. Math. Sci.]*, 38, Samara State Technical University, Samara, 2005, Pp.11–14. (In Russ.), <http://mi.mathnet.ru/eng/vsgtu/v38/p11>, DOI: <https://doi.org/10.14498/vsgtu363>. (Russ).
 15. Imanaliev M. I., Asanov A., Kadenova Z. A., A class of linear integral equations of the first kind with two independent variables.// *Doklady Mathematics*. – 2014. – Vol. 89. – No 1. – P. 98-102. – DOI 10.1134/S1064562414010281. – EDN XLBERD.
 16. Asanov A., M. H. Chelik, Kadenova Z. A., Uniqueness and stability of solutions of linear intergral equations of the first kind with two variables. // *International Journal of Mathematical Analysis*. – 2013. – Vol. 7. – No 17-20. – P. 907-914. – DOI 10.12988/ijma.2013.13088. – EDN XKWECX.
 17. Asanov A., Kadenova Z.A., Bekeshova D. On the uniqueness of solutions of Fredholm linear integral equations of the first kind on a semi-axis, // *Herald of Institute Mathematics of the National Academy of Sciences of the Kyrgyz Republic*. – 2022. – No 1. – P. 82-87. – DOI 10.52448/16948173_2022_1_82. – EDN FXALAA.
 18. Asanov A., Kadenova Z.A., Bekeshova D. On the uniqueness of solutions of Volterra linear equations of the first kind on the semi-axis// *Unstable problems of computational mathematics: Abstracts of the seminar with international participation, Irkutsk, August 15–19, 2022*. - Irkutsk: Irkutsk State University, 2022. - P. 36-40. – EDN KFMOZP.

Information about the authors

Asanov A. Doctor of Physical and Mathematical Sciences, Professor, Kyrgyz Turkish University Manas, Bishkek, ORCID iD: <https://orcid.org/0000-0002-5480-670X> e-mail: avyt.asanov@manas.edu.kg

Kadenova Z.A. Doctor of Physical and Mathematical Sciences, Institute of Mathematics of the National Academy of Sciences of the Kyrgyz Republic, ORCID iD: <https://orcid.org/0000-0002-3098-7446>, e-mail: kadenova71@mail.ru

Bekeshova D.A. Institute of Mathematics of the National Academy of Sciences of the Kyrgyz Republic, e-mail: bekeshova.d@mail.ru